


MODIFIED LIVE *FLAVOBACTERIUM COLUMNARE* AGAINST  
COLUMNARIS DISEASE IN FISH

Background of the Invention

Field of the Invention

[0001] *Flavobacterium columnare* is an aquatic bacterium that is highly infectious in both warm and cold water species of fish. In the channel catfish (*Ictalurus punctatus*), it is the causative agent of columnaris disease. *Flavobacterium columnare* is a Gram-negative, rod shaped, pathogen that has been isolated from channel catfish in areas of the southeastern United States where this species is cultured. The disease also affects sports fish (i.e., walleye and largemouth bass) and aquarium fishes. Medicated feed (antibiotics) is currently used to try and control this bacterial infection. However, these treatments are limited in their effectiveness and most producers have discontinued use of medicated feeds. Prevention of columnaris disease by vaccination is an important goal and a top priority of catfish

Express #	EK 419816454 US
I hereby certify that this correspondence is being deposited with the U.S. Postal Service as Express Mail in an envelope addressed to: Assistant Commissioner of Patents, Washington, D.C. 20231 on	
December 12, 2001 (Date of Deposit)	
Joseph A. Lipovsky Name of Depositor	
 Signature	
12/12/01 Date of Signature	

and other fish producers throughout the world. Estimated savings to these industries would be in excess of \$100 million annually.

[0002] This invention relates to a novel vaccine against columnaris which does in fact provide superior protection over commercial treatment involving treating the water with potassium permanganate ( $KmnO_4$ ) or feeding medicated feeds.

#### Description of the Prior Art

[0003] It has been estimated that columnaris is the second leading cause of mortality in pond raised catfish in the southeastern United States. Based on the success of disease control by immunization with killed bacteria (i.e., bacterins) in salmonids, experimental bacterins have been developed and tested against *F. columnare*. However, no vaccine is currently available and vaccination is not practiced in the catfish industry against *F. columnare*, presumably because the inactivation (i.e., formalin treatment) destroys the antigen (Bader et al., Comparison of whole-cell antigens of pressure-and formalin-killed *Flexibacter columnaris* from channel catfish (*Ictalurus punctatus*), American Journal of Veterinary Research, 58, pp 985-988, 1997). The patent of Wolf-Watz et al. (U.S. Patent 5,284,653) presents a whole list of bacteria of which one is *Flexibacter columnaris* (now *F. columnare*) which has the potential to be genetically modified to produce a vaccine. However, no data are presented on

*F. columnare* vaccines, only on genetically modified mutant vaccines of *Vibrio anguillarum*. Bernadet (Immunization with bacterial antigens: *Flavobacterium* and *Flexibacter* infections, Fish Vaccinology: Developments in Biological Standardization, Volume 90, pp 335-340, 1997; Karger Switzerland: Basel) reviews the limited knowledge available on *F. columnare* and vaccination against this important disease. Work suggests that rainbow trout (*Oncorhynchus mykiss*) which survive infection with *F. columnare* are immune to subsequent disease.

#### Summary of the Invention

[0004] We have now discovered a means for the creation of novel live vaccines that are safe and effective for the control of *F. columnare* in catfish. The vaccines comprise one or more rifampicin (3-[4-methylpiperazinyl-iminomethyl]rifamycin SV) (Sigma Chemical Company, St. Louis, MO) resistant mutants of *F. columnare*, created by multiple passaging of the native isolate on increasing concentrations of rifampicin. These vaccines are effective in providing long lasting acquired immunity in channel catfish to *F. columnare*.

[0005] In accordance with this discovery, it is an object of the invention to provide a novel, highly protective, live vaccine against *F. columnare* in fish, such as eels (*Anguilla* sp.), salmonids (*Oncorhynchus* sp. and *salmo* sp.), tilapia (*Oreochromis*

sp.), hybrid-striped bass (*Morone chrysops* x *M. saxatilis*), walleye (*Stizostedion vitreum*), channel catfish, cetrachids (such as largemouth bass (*Micropterus salmoides*)), bait minnows (*Pimephales promelas*), goldfish (*Carassius auratus*), carp (*Cyprinus carpio*), and aquarium fish (tropical fish species such as black mollies (*Poecilia sphenops*)) and platies (*Xiphophorus maculatus*).

[0006] It is another object to provide an attenuated *F. columnare* vaccine that is safe and provides long lasting acquired immunity in fish to columnaris disease, including channel catfish.

[0007] It is a further object of this invention to improve the viability and productivity of catfish, and to reduce economic losses in the fish industry caused by columnaris disease.

[0008] Other objects and advantages of the invention will become readily apparent from the ensuing description.

#### Deposit of Biological Material

[0009] Two rifampicin-resistant attenuated *F. columnare* isolate, were deposited on June 20, 2000 under the provisions of the Budapest Treaty in the Agricultural Research Service Culture Collection in Peoria, IL, and have been assigned Deposit No.'s B-30303 and B-30304, respectively.

## Detailed Description of the Invention

[0010] "Vaccine" is defined herein in its broad sense to refer to any type of biological agent in an administrable form capable of stimulating a protective immune response in an animal inoculated with the vaccine. For purposes of this invention, the vaccine may comprise one or more live attenuated mutants of *F. columnare* having the characteristic of rifampicin-resistance.

[0011] The starting material for use in preparing the vaccines of the invention is any attenuated *F. columnare* bacterium such as those reported supra. Serial passage of the isolate of *F. columnare* over increasing concentrations of rifampicin produces strains with an attenuated pathogenicity efficacious for the preparation of live vaccines. The attenuation achieved by high-level serial passage in culture on increasing concentrations of rifampicin virtually eliminates the pathogenicity of the bacterium toward fish. The native strain of *F. columnare* should be passaged a sufficient number of times such that in its new attenuated form it no longer possesses the ability of causing the disease state known as columnaris in catfish. The methodology for attenuation by serial passage is well known and documented in the art as exemplified by Schurig et al. [Vet. Micro. 28, 171-188 (1991)], hereby incorporated by reference, who created vaccines based on modified live rifampicin-resistant *Brucella* species.

[0012] Vaccination, while being accomplishable by injection or through oral ingestion, is most efficiently done by means of aqueous immersion. The bacterial agent is prepared for administration by formulation in an effective immunization dosage with an acceptable carrier or diluent, such as water. The expression "effective immunization dosage" is defined as being that amount which will induce immunity in a fish against challenge by a virulent strain of *Flavobacterium columnare*. Immunity is considered as having been induced in a population of fish when the level of protection for the population is significantly higher than that of an unvaccinated control group. One measure of protection following experimental challenge is relative percent survival (RPS) as described by Amend (1981; Dev. Biol. Stand., 49, 447-454), herein incorporated by reference. RPS is calculated according to the following formula:

$$RPS = 1 - \frac{\% \text{ vaccinee mortality}}{\% \text{ control mortality}} \times 100$$

[0013] A positive vaccinal effect is indicated by a RPS equal to or greater than 60%. Typically, vaccination is carried out by exposing fish by immersion in water containing about  $1 \times 10^6$  CFU/ml of attenuated *Flavobacterium columnare* for 15 minutes at a density of about 50 fish/L and a temperature of about 25°C. CFU denotes colony forming units of *F. columnare*. These parameters

may be varied as desired such that a sufficient level of vaccination is acquired without induction of stressful conditions or loss of fish. Useable concentrations of *Flavobacterium columnare* are considered to range from about  $5 \times 10^5$  to about  $1 \times 10^8$  CFU/ml of immersion medium. Useable vaccination times are seen to range from about 1 minute to about 60 minutes, preferably from about 2 minutes to about 15 minutes. Temperature of the inoculation media may range within the physiologically acceptable limits of the fish involved, for channel catfish preferably from about 18°C to about 28°C, most preferably from about 22°C to about 26°C. Concentrations of fish treated in the inoculation medium typically range from about 50 to about 100 fish/L, but, in the alternative, be determined on a weight basis and range from about 0.5 to about 2.5 kg/L. The vaccine can be effectively administered anytime after the fish attains immunocompetence, which for channel catfish is at about the second day to fourteen days post-hatch. Other species of fish susceptible to *F. columnare* can be immunized after 21-30 days post-hatch or when they become immunocompetent to modified live vaccine administered by immersion.

[0014] Appropriate adjuvants as known in the art may also be included in the vaccine formulation.

[0015] The following examples are intended only to further illustrate the invention and are not intended to limit the scope of the invention which is defined by the claims.

#### Example 1

[0016] The procedure used to produce the *F. columnare* vaccine mutants was modified from that described in Shurig et al. (1991; Vet Micro, 28, 171-188), hereby incorporated by reference, by using a lower initial concentration of rifampicin and ending at 200 µg/ml after 243 passages instead of 51 passages and by omitting the penicillin step.

#### Process of Developing Resistant Mutants of *Flavobacterium columnare*

[0017] Modified *Cytophaga* agar plates for the cultivation of *Flavobacterium columnare* were made according to the procedure of Klesius et al. (Effect of feed deprivation on innate resistance and antibody response to *Flavobacterium columnare* in channel catfish, *Ictalurus punctatus*. Bulletin European Association of Fish Pathologists, 19(4), 156-158, 1999). 1.0 g tryptone, 0.5 g yeast extract, 0.2 g beef extract, 0.2 g sodium acetate and 9.0 g of agar were added to one liter of distilled water. The media and agar were heated until dissolution. The media was then



autoclaved at 121-124°C for 15 minutes, the media was then poured into sterile petri dishes (15 ml per dish) and allowed to solidify before refrigeration.

[0018] Native isolates of *Flavobacterium columnare* were obtained from sick catfish or previously obtained lyophilized stocks. Isolates of *F. columnare* were then identified by standard biochemical tests as set forth in Bergey's Manual of Determinative Bacteriology prior to use in rifampicin resistant *F. columnare*. After identification, the process of forming rifampicin resistant isolates of *F. columnare* was begun. Rifampicin supplemented modified *Cytophaga* agar plates were prepared as follows: Modified *Cytophaga* agar was made as described above and sterilized at 121-124°C for 15 minutes. After sterilization, the correct amount of rifampicin was added to the media prior to its solidification and 15 ml of the resulting mixture was poured into separate petri dishes and allowed to solidify prior to refrigerated storage. Initial cultures of the native isolates of *F. columnare* were grown on modified *Cytophaga* agar plates which were incubated at 20-25°C for 24-48 hours or until 1-2 mm yellow, rhizoid colonies were observed. A single *F. columnare* colony was then picked with a sterile inoculating loop and streaked onto a rifampicin supplemented modified *Cytophaga* agar plate containing the correct concentration of the antibiotic. For the initial passage,

rifampicin was present in the modified *Cytophaga* agar at a concentration of 5  $\mu\text{g/ml}$ . The rifampicin supplemented modified *Cytophaga* agar which was streaked with the aforementioned native isolate of *F. columnare* was then incubated for 24-48 hours at 20-25°C and observed for bacterial growth. Single colonies of *F. columnare* which grew on the rifampicin supplemented media were then picked and placed onto the next concentration of rifampicin (10  $\mu\text{g/ml}$ ) modified *Cytophaga* agar plates. If growth occurred, a single colony was harvested and placed on an agar media containing the next higher concentration of rifampicin (20  $\mu\text{g/ml}$ ). If the harvested colony failed to grow, it was repeatedly passed on a media containing the last concentration of rifampicin at which growth successfully occurred, before being placed on the next higher concentration of rifampicin containing media. This process was repeated until a colony capable of growing on a media containing a rifampicin concentration of 200  $\mu\text{g/ml}$  was created.

[0019] *Flavobacterium columnare* isolate ARS-1 was passaged on increasing concentrations of rifampicin (Sigma Chemical Company, St. Louis, MO) supplemented modified *Cytophaga* agar to a final concentration of 200  $\mu\text{g/ml}$  rifampicin for 243 passages. The resultant mutants (i.e., two colonies from the original passage that grew and were passaged), designated B-30303 and B-30304, are differentiated from the parent microorganism because they can

survive and reproduce on a media containing 200 µg/ml rifampicin without negative effect. Biochemical characteristics of the *F. columnare* B-30303 and B-30304 are identical to *F. columnare* as described in Bergey's Manual of Determinative Bacteriology (Holt et al., 1994), herein incorporated by reference.

## Example 2

### SAFETY AND BACK-PASSAGE

[0020] The safety and back-passage study revealed both vaccine candidate rifampicin mutants were safe for use in vaccination of 10 day post hatch channel catfish at concentrations of  $1.4 \times 10^8$  or  $6.7 \times 10^7$  CFU/ml for a 30 minute immersion exposure. One hundred and twenty five fish were used for each group which included the initial exposure and then each subsequent exposure for a total of 5 passages (i.e., fish to fish transfer). One hundred channel catfish not exposed to *F. columnare* mutants were kept as negative controls. The numbers of fish which died after exposure to the vaccine or vaccine diluent (i.e., controls) are presented in Table 1. Forty eight hours following exposure, 40 fish were removed, homogenized and cultured for the presence of *F. columnare* vaccine strains at each passage and in the control group. The vaccine isolates were isolated at the first passage but not at subsequent passages indicating that the vaccine strains were capable of invading the fish, but did not revert to

virulence following back-passage and thus, were attenuated in 10 day post hatch channel catfish. The fish used in this experiment were held in the laboratory without signs of columnaris disease or adverse behavior for at least 21 days following treatment.

TABLE 1

SAFETY OF VACCINE DOSE OF *F. COLUMNARE* MUTANTS IN  
10 DAY POST HATCH CHANNEL CATFISH VACCINATES

Mutant	Passage	No. Dead/ No. Total <sup>1</sup>	<i>F. columnare</i> Mutant Isolated from homogenized fish	Percent Mortality
B-30303	1	14 <sup>2</sup> /125	Yes	11.2
	2	0/125	No	0.0
	3	0/125	No	0.0
	4	0/125	No	0.0
	5	0/125	No	0.0
B-30304	1	17 <sup>3</sup> /125	Yes	13.6
	2	1 <sup>4</sup> /125	No	8.0
	3	12 <sup>5</sup> /125	No	9.6
	4	2 <sup>6</sup> /125	No	1.6
	5	0/125	No	0.0
Controls	None	13 <sup>7</sup> /100	No	13.0

<sup>1</sup>Note, total number of fish to start the experiment was used for calculations.

<sup>2</sup>1 of 14 positive for rifampicin resistant *F. columnare*.

<sup>3</sup>1 of 17 positive for rifampicin resistant *F. columnare*.

<sup>4</sup>*F. columnare* not isolated from dead fish.

<sup>5</sup>*F. columnare* not isolated from dead fish.

<sup>6</sup>*F. columnare* not isolated from dead fish.

<sup>7</sup>*F. columnare* not isolated from dead fish.

### EXAMPLE 3

#### Efficacy

[0021] Channel catfish (125, USDA 103 strain) were vaccinated by adding 40 ml of *F. columnare* mutant B-30303 to 7.5 L of water for a 30 minute exposure. One-hundred twenty five USDA 103 channel catfish were exposed by adding 40 ml of vaccine diluent to 7.5 L of water for a 30 minute exposure to serve as control fish (i.e., non-vaccinated). Vaccinated and control fish were held for 32 days following vaccination before they were challenged with virulent *F. columnare*. No fish died after the vaccination. Results of experimental challenge are presented as relative percent survival (RPS) as described by Amend (1981; Dev. Biol. Stand., 49, 447-454), herein incorporated by reference. RPS is calculated according to the following formula:

$$RPS = 1 - \frac{\% \text{ vaccinate mortality}}{\% \text{ control mortality}} \times 100$$

[0022] A positive effect by vaccination is a RPS greater than 50%. At fourteen days post vaccination (DPV), the relative percent survival (RPS) was 69.2%. In this study, mortality in an equivalent group of 125 untreated controls was 20.8% versus 6.4% mortality in the 125 vaccinated fish.

TABLE 2

PROTECTION AGAINST COLUMNARIS DISEASE AFTER IMMERSION  
VACCINATION<sup>1</sup> OF CHANNEL CATFISH  
WITH *FLAVOBACTERIUM COLUMNARE* B-30303 VACCINE

Treatment	No. Dead/ No. Total	Percent Mortality	Relative Percent Survival (RPS) <sup>2</sup>
Vaccinated with B-30303	8/125	6.4	69.2
Control (non-vaccinated)	26/125	20.8	--

<sup>1</sup>Immersion vaccination for 30 minutes with  $5 \times 10^6$  CFU/ml  
*F. columnare* B-30303 for 30 minute immersion exposure.

<sup>2</sup>Relative percent survival as determined by Amend (1981).

## EXAMPLE 4

[0023] Fish were vaccinated (48 day post hatch channel catfish) with the modified live *F. columnare* B-30303 for two minutes at the described vaccine dose and then diluted with an equal volume of water and held for an additional 13 minutes (total vaccine time 15 minutes). Control and control non-challenged groups were immersed in vaccine diluent for the same duration of exposure (15 minutes). The fish were challenged 57 days following vaccination by co-habitation following 14 days without feed (Klesius et al., 1999). Three channel catfish showing signs of columnaris disease were added to each of the groups A, B and C and allowed to remain with fish for 24 hours. A sub-sample of five dead fish of those used for cohabitation

were culture positive for *F. columnare*. The dead fish were removed at 24 hours the following day and feeding resumed. Table 3 shows the relative percent survival was calculated according to Amend (1981).

Table 3

PROTECTION AGAINST COLUMNARIS DISEASE AFTER  
IMMERSION VACCINATION OF CHANNEL CATFISH WITH  
*FLAVOBACTERIUM COLUMNARE* B-30303 VACCINE

Group	Treatment	No. Dead/ No. Total	Mean % Mortality (SEM <sup>1</sup> )	Relative Percent Survival <sup>2</sup>
A	Vaccinated 1 x 10 <sup>6</sup> CFU/ml	2/150	1.33 (0.67) <sup>a</sup>	96.4
B	Vaccinated 5 x 10 <sup>6</sup> CFU/ml	7/150	4.67 (0.67) <sup>a</sup>	97.3
C	Control	55/150	36.67 (17.7) <sup>b</sup>	---
D	Control Non-challenged	3/150	2.00 (1.15) <sup>a</sup>	---

<sup>1</sup>Means with different superscripts are significantly different at  $p < 0.05$  using Duncan's Multiple range test for differences (SAS Inc., 1997).

<sup>2</sup>Relative percent survival calculated according to Amend (1981).

EXAMPLE 5

[0024] *Flavobacterium columnare* isolates were obtained from channel catfish with signs of columnaris disease. The *F. columnare* were cultured on *Cytophaga* agar and determined to be pure cultures. Isolates obtained were frozen in 2 ml aliquots at

-80°C. Once thawed to 25°C, fifty microliters were then plated onto each of the media types (i.e., *Cytophaga* agar and *Cytophaga* agar supplemented with 5 µg rifampicin/ml) and incubated at 25±3°C for 24 hours. The development of yellow rhizoid colonies on agar was considered positive for growth. The ten *F. columnare* isolates tested grew on *Cytophaga* agar only with typical *F. columnare* colonies developing. Growth was not observed with any of the isolates tested on *Cytophaga* agar supplemented with 5 µg rifampicin/ml.

[0025] Culture was also attempted after the isolates recovered from cold storage had grown on the non-rifampicin supplemented plates for 48 hours. Single colonies of the recovered bacteria that were growing on *Cytophaga* agar were picked and streaked onto *Cytophaga* agar supplemented with 5 µg rifampicin/ml. No growth occurred on *Cytophaga* agar supplemented with 5 µg rifampicin/ml following the use of 48 hours cultures of recovered *F. columnare*. Lack of growth on *Cytophaga* agar supplemented with 5 µg rifampicin/ml demonstrated that these isolates were not viable candidates for development of modified live *F. columnare* vaccines because rifampicin resistance could not be induced in these native *F. columnare* isolates (see Table 4). Failure of induction of rifampicin resistance was demonstrated by no growth on rifampicin supplemented *Cytophaga* agar. This testing shows the unexpected nature of the success achieved in Example 1.



Table 4

Failure of ten *Flavobacterium columnare* isolates  
tested to grow on rifampicin supplemented  
Cytophaga agar (5 µg rifampicin/ml)

Isolate Identification <sup>1</sup>	Growth on Cytophaga Agar	Growth on Cytophaga rifampicin supplemented agar (5 µg rifampicin/ml)
AUFAA-1 <sup>2</sup>	Yes	No Growth
AUFAA-2	Yes	No Growth
AUFAA-3	Yes	No Growth
AUFAA-4	Yes	No Growth
AUFAA-5	Yes	No Growth
ALG-530 <sup>3</sup>	Yes	No Growth
ALG-515	Yes	No Growth
ALG-513	Yes	No Growth
ALG-527	Yes	No Growth
ALG-521	Yes	No Growth

<sup>1</sup>*Flavobacterium columnare* isolates all from channel catfish  
showing signs of columnaris disease.

<sup>2</sup>AUFAA = Auburn University Fisheries and Allied Aquaculture -  
Isolate Number.

<sup>3</sup>ALG = Alabama Fish Farming Center, Greensboro, AL - Case Number.

[0026] It is understood that the foregoing detailed  
description is given merely by way of illustration and that  
modification and variations may be made therein without departing  
from the spirit and scope of the invention.